

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

aTD224
.N2N48

United States
Department of
Agriculture

Soil
Conservation
Service

Reno
Nevada



STA/ST

Nevada Water Supply Outlook

January 1, 1986



Foreward

How Forecasts Are Made

Most of the annual streamflow in the Western United States originates as snowfall. This snowfall accumulates high in the mountains during winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Predictions are based on careful measurements of snow water equivalent at selected index points. Precipitation, temperature, soil moisture and antecedent streamflow data are viewed in conjunction with snowpack data to prepare runoff forecasts. This report presents a comprehensive picture of water supply outlook conditions for areas dependent upon surface runoff. It includes selected streamflow forecasts, summarized snowpack and precipitation data, reservoir storage data and narratives describing current conditions.

Streamflow forecasts are cooperatively generated by Soil Conservation Service and National Weather Service hydrologists. Forecasts become more accurate as more data affecting runoff becomes known. For this reason, forecasts are issued that reflect three future precipitation conditions — Below Normal, Average, and Above Normal. These forecasts are termed reasonable minimum, most probable, and reasonable maximum. Actual streamflow can be expected to fall between the lower and upper forecast values eight out of ten years.

Snowpack data are obtained by using a combination of manual and automated measurement methods. Manual readings of snow depth and water equivalent are taken at locations called snow courses on a monthly or semi-monthly schedule during the winter. In addition, snow water equivalent, precipitation, temperature, and other parameters are monitored on a daily basis and transmitted via radio telemetry to central data collection facilities. Both monthly and daily data are used to project snowmelt runoff.

For More Information

Copies of Monthly Water Supply Outlook Reports and other reports may be obtained from the states listed below. Because of the limited space, snow survey measurements are not published in monthly reports. An annual snow survey data summary is published by the Soil Conservation Service for each of the western states. Historical snow survey data may be obtained at those same offices.

STATE	ADDRESS
Alaska	201 East 9th Ave., Suite 300, Anchorage, AK 99501-3687
Arizona	201 East Indianola, Suite 200, Phoenix, AZ 85012
Colorado (New Mexico)	2490 West 26th Ave., Denver, CO 80211
Idaho	304 North 8th Street, Room 345, Boise, ID 83702
Montana	10 East Babcock, Room 443, Federal Building, Bozeman, MT 59715
Nevada	1201 Terminal Way, Second Floor, Reno, NV 89502
Oregon	1220 Southwest 3rd Ave., 16th Floor, Portland, OR 97204
Utah	4402 Federal Building, 125 South State Street, Salt Lake City, UT 84147
Washington	360 U.S. Court House, Spokane, WA 99201
Wyoming	Federal Building, 100 East "B" Street, Casper, WY 82602

In addition to state reports, a Water Supply Outlook for the Western United States is published by the Soil Conservation Service and National Weather Service monthly, January through May. Reports may be obtained from the Soil Conservation Service, West National Technical Center, 511 Northwest Broadway, Room 547, Portland, OR 97209.

Published by other agencies:

Water Supply Outlook Reports prepared by other agencies include: California — Snow Survey Branch, California Department of Water Resources, P.O. Box 388, Sacramento, CA 98502; British Columbia — The Ministry of Environment, Water Investigations Branch, Parliament Buildings, Victoria, British Columbia, V8V 1X5; Yukon Territory — Department of Indian and Northern Affairs, Northern Operations Branch, 200 Range Road, Whitehorse, Yukon Territory, Y1A 3V1; Alberta, Saskatchewan, and N.W.T. — The Water Survey of Canada, Inland Waters Branch, 110-12 Avenue S.W., Calgary, Alberta, T3C 1A6.

Nevada Water Supply Outlook

and

Federal - State - Private Cooperative Snow Surveys

Issued By

Wilson Scaling
Chief
Soil Conservation Service
Washington, DC 20013

Released By


Gerald Thola
State Conservationist
Soil Conservation Service
Reno, Nevada 89502

Prepared By

John R. Capurro
Water Supply Specialist
Soil Conservation Service
1201 Terminal Way, Second Floor
Reno, Nevada 89502

In Cooperation With

Roland D. Westergard
Director
Department of Conservation &
Natural Resources
Carson City, Nevada 89701



GENERAL OUTLOOK

SUMMARY:

WESTERN NEVADA SNOWPACK ACCUMULATIONS ARE AVERAGE WHILE NORTHERN AND EASTERN NEVADA ARE SIGNIFICANTLY ABOVE AVERAGE. PRECIPITATION AT MOST SNOTEL SITES WAS BELOW AVERAGE FOR DECEMBER, BUT IS NEAR AVERAGE FOR THE WATER YEAR. RESERVOIR STORAGE IN THE SEVEN MONITORED RESERVOIRS AND LAKES IS NEAR AVERAGE.

SNOWPACK:

Snowpack conditions in the Tahoe, Truckee, Carson, and Walker basins are average as of January 1. Snow accumulations in northern and eastern Nevada are slightly above to significantly above average. The Snake Basin is 10-15 percent above January 1 averages while the Owyhee, Humboldt and Eastern Nevada basins are much above average. The Northern Nevada basin is approximately 10 percent above average.

PRECIPITATION:

December precipitation totals ranged from slightly below average to much above average. The Snake and Owyhee basins were near average for the month. Water year accumulations in the Tahoe-Truckee basin is near average while all other basins are above to much above average. The Northern Great and Humboldt basins are approximately 120 percent of January 1 averages. The Carson-Walker basin is 115 percent of average.

RESERVOIRS:


Water storage is good throughout the state. Storage facilities in the Tahoe-Truckee, Humboldt, and Snake-Owyhee basins are well above average for January 1. Storage in the Carson-Walker basin is approximately 15 percent below average. Total storage in the seven major lakes and reservoirs was 792,000 acre feet.

STREAMFLOW:

Streamflow forecasts for western Nevada are near or slightly below average. Forecast values for all northern Nevada rivers and creeks are much above average. Streamflow forecast values range from 5 percent below average for the Carson River at Carson City to 35 percent above average for the Franklin River near Arthur, Nevada.

TAHOE & TRUCKEE BASINS

Mountain snowpack* (inches)




*Based on selected stations

Maximum
Minimum

Average
Current

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation Year to date precipitation

WATER SUPPLY OUTLOOK:

Snowpack accumulations are near normal. This year's water content is approximately 10 percent below the values recorded last year. Reservoir storage is excellent with all storage facilities in the basin at or well above average. Storage in Boca, Prosser, and Stampede reservoirs is less than the amount stored last year at this date. Streamflow forecasts are average to slightly above average for Truckee River gaging stations.

For more information contact your local Soil Conservation Service office.

TAHOE & TRUCKEE BASINS

STREAMFLOW FORECASTS


FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
LAKE TAHOE RISE (assume gates closed APR-HIG)		1.3	1.4	100.0	215.8	0.0				
TRUCKEE RIVER at Farad, Ca	APR-JUL	269.0	265.0	98.0	181.4	19.7				
LITTLE TRUCKEE RIVER above Boca, Ca	APR-JUL	92.5	97.0	104.0	189.2	34.6				
PYRAMID LAKE RISE	OCT-HIG	1.0	-0.2		100.0	8.0				
STEAMBOAT CREEK at Steamboat, Nv	APR-JUL	5.2	4.8	92.0	173.1	19.2				
SAGEHEN CREEK, Ca	APR-JUL	6.5	6.2	95.0	184.6	15.4				
GALENA CREEK nr Steamboat, Nv	APR-JUL	4.4	4.3	97.0	181.8	22.7				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	USEABLE STORAGE THIS YEAR	USEABLE STORAGE LAST YEAR	USEABLE STORAGE AVE.	WATERSHED	NO. COURSES AVE.D	THIS YEAR AS % OF LAST YR.	% OF AVERAGE
BOCA RESERVOIR	40.9	22.3	23.0	17.0	LAKE TAHOE RISE	8	93	107
LAKE TAHOE	744.6	428.2	543.0	359.2	TRUCKEE BASIN	8	102	108
PROSSER RESERVOIR	28.6	9.5	9.0	7.5	LITTLE TRUCKEE RIVER	3	105	108
STAMPEDE RESERVOIR	226.5	118.3	192.0	95.6	SAGE HEN CREEK	3	105	108
					GALENA CREEK	3	99	104
					STEAMBOAT DRAINAGE	2	98	98
					PYRAMID LAKE	16	98	108

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

CARSON & WALKER BASINS


Mountain snowpack* (inches)



*Based on selected stations

Maximum ——— Average - - - -
Minimum ——— Current ———

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation  Year to date precipitation 

WATER SUPPLY OUTLOOK:

Water content values are average for January 1. Reservoir storage is 10-15 percent below average for the month. Bridgeport and Topaz reservoirs in the Walker drainage are 15 percent below average while Lahontan reservoir in the Carson drainage is 10 percent below average. Bridgeport and Lahontan reservoirs have less stored water this year than last while Topaz Lake has slightly more stored water. Streamflow forecasts are average to slightly below average.

For more information contact your local Soil Conservation Service office.

CARSON & WALKER BASINS

STREAMFLOW FORECASTS


FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
E.F. CARSON RIVER near Gardnerville, Nv	APR-JUL	187.0	197.0	105.0	144.0	65.0	1846		200	JUN 21
W.F. CARSON RIVER at Woodfords, Ca	APR-JUL	53.0	54.0	101.0	139.6	66.0				
CARSON RIVER near Carson City, Nv	APR-JUL	182.0	186.0	102.0	172.0	32.4	1900			
CARSON RIVER near Ft. Churchill, Nv	APR-JUL	166.0	169.0	101.0	184.9	18.7	1837			
EAST WALKER RIVER near Bridgeport, C	APR-JUL	66.0	68.0	103.0	169.7	36.4				
WEST WALKER RIVER near Coleville, Ca	APR-JUL	148.0	155.0	104.0	157.4	52.0	1573			
WALKER LAKE RISE	OCT-HIG	1.0	-0.3		100.0	3.0				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	THIS YEAR	LAST YEAR	AVE.	WATERSHED	NO. COURSES AVE.D	THIS YEAR AS % OF LAST YR.	% OF AVERAGE
BRIDGEPORT RESERVOIR	42.5	18.7	30.0	23.7	E. CARSON RIVER	5	74	102
LAHONTAN RESERVOIR	295.1	145.5	148.0	165.6	W. CARSON RIVER	3	76	98
TOPAZ RESERVOIR	59.4	20.6	13.0	26.9	CARSON RIVER at Carson Ci	4	71	100
					CARSON RIVER at Ft. Churc	4	71	100
					E. WALKER RIVER nr Bridge	3	77	102
					W. WALKER RIVER nr Colevi	4	84	101
					WALKER LAKE RISE	4	84	101

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

HUMBOLDT BASIN


Mountain snowpack* (inches)





*Based on selected stations

Maximum ——— Average - - - - -
Minimum = = = Current ———

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation  Year to date precipitation 

WATER SUPPLY OUTLOOK:

Snowpack accumulations are significantly above average for the fourth consecutive year. The January 1 snowpack is approximately 200 percent of average. Basin water year precipitation is above average by about 10 percent. Storage in Rye Patch Reservoir is approximately 30 percent above average for this date. Streamflow forecasts for the Humboldt River are 185 percent of average.

For more information contact your local Soil Conservation Service office.

HUMBOLDT BASIN

STREAMFLOW FORECASTS

FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
HUMBOLDT RIVER at Palisades	APR-JUL	230.0	275.0	119.0	212.6	26.5				
HUMBOLDT RIVER at Comus	APR-JUL	173.0	202.0	116.0	239.9	15.6				
S FORK HUMBOLDT RIVER at Dixie	APR-JUL	75.0	88.0	117.0	201.3	33.3				
NF HUMBOLDT RIVER at Devils Gate	APR-JUL	34.8	40.0	114.0	215.5	14.4				
MARY'S RIVER nr Deeth	APR-JUL	36.9	40.0	108.0	173.4	43.4				
MARTIN CREEK nr Paradise Nv	APR-JUL	15.8	17.4	110.0	177.2	38.0				
LAMOILLE CREEK nr Lamoille	APR-JUL	28.7	29.0	101.0	149.8	52.3				
REESE RIVER nr Ione Nv	APR-JUL	6.6	7.0	106.0	197.0	15.2				
LITTLE HUMBOLDT RIVER nr Paradise Va	APR-JUL	9.7	11.2	115.0	185.6	51.5				
ROCK CREEK nr Battle Mtn.	APR-JUL	16.0	17.6	110.0	193.7	25.0				
KINGSTON CREEK nr Austin, Nv	APR-JUL	3.3	3.9	118.0	212.1	30.3				
MCDERMITT CREEK nr McDermitt	APR-JUL	12.0	14.0	116.0	183.3	50.0				

RESERVOIR STORAGE


(1000AF)

WATERSHED SNOWPACK ANALYSIS

RESERVOIR	USEABLE CAPACITY	USEABLE STORAGE THIS YEAR	USEABLE STORAGE LAST YEAR	USEABLE STORAGE AVE.	WATERSHED	NO. COURSES AVE.D	THIS YEAR AS % OF LAST YR.	% OF AVERAGE
RYE PATCH RESERVOIR	194.3	118.5	158.0	88.7	LAMOILLE CREEK	1	105	226
					S. FORK HUMBOLDT	4	109	171
					MARY'S RIVER	3	75	112
					N. FORK HUMBOLDT	3	167	232
					HUMBOLDT RIVER at Palisad	6	116	187
					HUMBOLDT RIVER at Comus	6	116	187
					KINGSTON CREEK	1	89	192
					LITTLE HUMBOLDT RIVER	1	123	286

SNAKE & Owyhee Basins


Mountain snowpack* (inches)



*Based on selected stations

Maximum ——— Average - - - - -
Minimum ——— Current ———

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation Year to date precipitation

WATER SUPPLY OUTLOOK:

The accumulated snowpack is well above average for January 1 with water content values as much as 65 percent above 20 year averages and 5 percent above last year's values. Water year precipitation is also above average. Wildhorse Reservoir is approximately 40 percent above average for this time of year, but is well below the amount stored last year. Streamflow forecasts for the three gauging stations within the basins are 110 percent of average.

For more information contact your local Soil Conservation Service office.

SNAKE & OWYHEE BASINS

STREAMFLOW FORECASTS


FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
OWYHEE RIVER nr Gold Creek	APR-JUL	23.4	26.0	111.0	188.0	34.2				
OWYHEE RIVER nr Owyhee	APR-JUL	85.4	96.0	112.0	185.0	39.8				
S FORK OWYHEE nr White Rock, Nv	APR-JUL	83.0	90.0	108.0	181.9	34.9				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	USEABLE STORAGE THIS YEAR	USEABLE STORAGE LAST YEAR	USEABLE STORAGE AVE.	WATERSHED	NO. COURSES AVER.D	THIS YEAR AS % OF LAST YR.	% OF AVERAGE
MILDHORSE RESERVOIR	71.5	38.2	60.0	26.6	OWYHEE RIVER nr Owyhee	4	106	165
					OWYHEE RIVER nr Gold Cree	1	166	166
					S. FORK OWYHEE RIVER	4	106	165
					SALMON FALLS CREEK	3	75	112

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

EASTERN NEVADA


Mountain snowpack* (inches)



*Based on selected stations

Maximum ——— Average - - - - -
Minimum ——— Current ———

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation Year to date precipitation

WATER SUPPLY OUTLOOK:

Snowpack water content accumulation is well above average for January 1. The current water content is approximately equal to the amounts measured last year at this time. Streamflow forecasts for Steptoe Creek and Franklin River are above average by 20 percent and 35 percent respectively.

For more information contact your local Soil Conservation Service office.

EASTERN NEVADA

STREAMFLOW FORECASTS


FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
STEPTOE CREEK nr Elv	APR-JUL	2.0	2.4	119.0	200.0	50.0				
FRANKLIN RIVER nr Arthur	APR-JUL	5.9	8.0	135.0	220.3	50.8				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	USEABLE THIS YEAR	STORAGE LAST YEAR	USEABLE STORAGE AVERAGE	WATERSHED	NO. COURSES AVERAGE	THIS YEAR LAST YR.	AS % OF AVERAGE
					FRANKLIN RIVER	1	110	155
					EASTERN NEVADA	1	88	182
					STEPTOE VALLEY	0	0	0

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

NORTHERN GREAT BASIN


Mountain snowpack* (inches)



*Based on selected stations

Maximum ——— Average - - - - -
 Minimum ——— Current ———

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation Year to date precipitation

WATER SUPPLY OUTLOOK:

Snowpack accumulations in this basin are variable. Water content values in the western and southern portions are below average while water content in the eastern and northern portions is above average. All water content values are below those reported last year. Streamflow forecasts for the Quinn River, East Fork Quinn River, and McDermitt Creek are approximately 15 percent above average. Streamflow forecast values for Deep Creek, Eagle Creek, and Mill Creek are approximately 10 percent below average.

For more information contact your local Soil Conservation Service office.

NORTHERN GREAT BASIN

STREAMFLOW FORECASTS


FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
BIDWELL CREEK nr Fort Bidwell	APR-JUL	12.0	11.3	94.0	175.0	16.7				
DEEP CREEK nr Cedarville, Ca	APR-JUL	3.6	3.2	88.0	166.7	0.0				
EAGLE CREEK nr Eagleville, Ca	APR-JUL	4.3	4.1	95.0	162.8	23.3				
MILL CREEK nr Cedarville, Ca	APR-JUL	4.1	3.8	92.0	170.7	24.4				
QUINN RIVER nr McDermitt, Nv	APR-JUL	16.0	16.8	105.0	175.0	37.5				
E. FORK QUINN RIVER nr McDermitt	APR-JUL	13.0	14.0	107.0	176.9	38.5				
MCDERMITT CREEK nr McDermitt	APR-JUL	12.0	14.0	116.0	183.3	50.0				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	** USEABLE STORAGE **			WATERSHED	NO. COURSES AVE.D	THIS YEAR AS % OF	
		THIS YEAR	LAST YEAR	AVE.			LAST YR.	AVERAGE
					BIDWELL	1	47	70
					MILL CREEK	1	47	70
					DEEP CREEK	1	47	70
					EAGLE CREEK	1	47	70
					QUINN RIVER	2	69	116
					E. FORK QUINN	2	69	116
					MCDERMITT CREEK	2	69	116

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

SOUTHERN NEVADA


Mountain snowpack* (inches)



*Based on selected stations

Maximum Average
 Minimum Current

Precipitation* (percent of normal)



*Based on selected stations

Monthly precipitation Year to date precipitation

WATER SUPPLY OUTLOOK:

Snowpack accumulations in the mountains supplying water for the Virgin River are approximately 15 percent above average for January 1. Storage in Lake Mohave is about 10 percent below average while storage in Lake Mead is about 30 percent above average. Both storage values are less than those recorded last year at this time. The streamflow forecast for the Virgin River near Hurricane, Utah is 85 percent above the 20 year average.

For more information contact your local Soil Conservation Service office.

SOUTHERN NEVADA

STREAMFLOW FORECASTS

FORECAST POINT	FORECAST PERIOD	20 YR. AVE. (1000AF)	MOST PROBABLE (1000AF)	MOST PROBABLE (% AVE.)	REAS. MAX. (% AVE.)	REAS. MIN. (% AVE.)	PEAK FLOW (CFS)	PEAK DATE	LOW FLOW (CFS)	LOW DATE
VIRGIN RIVER near Hurricane, UT	APR-JUL	48.0	90.0	187.0	252.1	145.8				

RESERVOIR STORAGE (1000AF)					WATERSHED SNOWPACK ANALYSIS			
RESERVOIR	USEABLE CAPACITY	USEABLE THIS YEAR	USEABLE STORAGE LAST YEAR	USEABLE STORAGE AVE.	WATERSHED	NO. COURSES AVE.D	THIS YEAR AS % OF LAST YR.	% OF AVERAGE
LAKE MOHAVE	1810.0	1422.0	1523.0	1605.0	VIRGIN RIVER at Littlefie	3	60	116
LAKE MEAD	26159.0	23683.0	24070.0	18176.0	VIRGIN RIVER at Hurricane	3	60	116

*Corrected for upstream diversions or changes in reservoir storage.
Average is for 1961-80 period.

The Following Organizations Cooperate With The Soil Conservation Service In Snow Survey Work

STATE

California Cooperative Snow Surveys
California Department of Parks and Recreation
California Department of Water Resources
Colorado River Commission of Nevada
Idaho Cooperative Snow Surveys
Nevada Association of Conservation Districts
Nevada Department of Conservation & Natural Resources
 Division of Water Resources
 Nevada State Forester
 Division of Conservation Districts
Oregon Cooperative Snow Surveys
University of Nevada, Desert Research Institute
Utah Cooperative Snow Surveys

FEDERAL

Bureau of Reclamation
Forest Service
Geological Survey
Soil Conservation Service
U.S. District Court - Federal Water Master
NOAA, National Weather Service

PRIVATE

Nevada Irrigation District
Owyhee Project North Board of Control
Owyhee Project South Board of Control
Pacific Gas and Electric Company
Pershing County Water Conservation District
Sierra Pacific Power Company
Truckee - Carson Irrigation District
Walker River Irrigation District
Washoe County Water Conservancy District

Other organizations and individuals furnish valuable information for the snow survey reports. Their cooperation is gratefully acknowledged.

UNITED STATES DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
1201 TERMINAL WAY, SECOND FLOOR
RENO, NEVADA 89502

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

THIRD CLASS BULK RATE
POSTAGE AND FEES PAID
USDA • SCS
RENO, NV
PERMIT NO G-3851

THIRD CLASS MAIL

Nevada
Water Supply Outlook
and

Federal — State — Private
Cooperative Snow Surveys



SOIL CONSERVATION SERVICE